

The Conservative Management of Idiopathic Toe Walking Utilizing an Evidence-Based Algorithm and Sensory-Motor Treatment Approach: References¹⁻⁹²

1. Alanen JT, Levola JV, Helenius HY, Kvist MH. Ankle joint complex mobility of children 7 to 14 years old. *J Pediatr Orthop.* Nov-Dec 2001;21(6):731-737.
2. Alvarez C, De Vera M, Beauchamp R, Ward V, Black A. Classification of idiopathic toe walking based on gait analysis: development and application of the ITW severity classification. *Gait Posture.* Sep 2007;26(3):428-435.
3. Armand S, Watelain E, Mercier M, Lensed G, Lepoutre FX. Identification and classification of toe-walkers based on ankle kinematics, using a data-mining method. *Gait Posture.* Feb 2006;23(2):240-248.
4. Ayres A. *Sensory integration and learning disorders.* Los Angeles: Western Psychological Services; 1972.
5. Bandy WD, Irion JM. The effect of time on static stretch on the flexibility of the hamstring muscles. *Phys Ther.* Sep 1994;74(9):845-850; discussion 850-842.
6. Bandy WD, Irion JM, Briggler M. The effect of time and frequency of static stretching on flexibility of the hamstring muscles. *Phys Ther.* Oct 1997;77(10):1090-1096.
7. Bandy WD, Irion JM, Briggler M. The effect of static stretch and dynamic range of motion training on the flexibility of the hamstring muscles. *J Orthop Sports Phys Ther.* Apr 1998;27(4):295-300.
8. Bennell KL, Talbot RC, Wajswelner H, Techovanich W, Kelly DH, Hall AJ. Intra-rater and inter-rater reliability of a weight-bearing lunge measure of ankle dorsiflexion. *Aust J Physiother.* 1998;44(3):175-180.
9. Brouwer B, Davidson LK, Olney SJ. Serial casting in idiopathic toe-walkers and children with spastic cerebral palsy. *J Pediatr Orthop.* Mar-Apr 2000;20(2):221-225.
10. Campbell S, Palisano R, Orlin M. *Physical Therapy for Children.* 4th ed: Saunders, an imprint of Elsevier, Inc.; 2012.
11. Chard A, Greene A, Hunt A, Vanwanseele B, Smith R. Effect of thong style flip-flops on children's barefoot walking and jogging kinematics. *J Foot Ankle Res.* 2013;6(1):8.
12. Clark E, Sweeney JK, Yocom A, McCoy SW. Effects of motor control intervention for children with idiopathic toe walking: a 5-case series. *Pediatr Phys Ther.* Winter 2010;22(4):417-426.
13. Coledam D, de Arruda G, de Oliveira A. Chronic effect of static stretching performed during warm-up on flexibility in children. *Brazilian Journal of Kinanthropometry and Human Performance.* 2012;14(3):296-304.
14. Conrad L, Bleck EE. Augmented auditory feed back in the treatment of equinus gait in children. *Dev Med Child Neurol.* Dec 1980;22(6):713-718.
15. Crenna P, Fedrizzi E, Andreucci E, Frigo C, Bonob R. The heel-contact gait pattern of habitual toe walkers. *Gait and Posture.* 2005;21:311-317.
16. Curry J, Exner C. Comparison of tactile preferences in children with and without cerebral palsy. *Am J Occup Ther.* Jun 1988;42(6):371-377.
17. Cusick B. *Serial casting for the restoration of soft-tissue extensibility in the ankle and foot: Scientific rationale and clinical management principles.* Progressive Gaitways, LLC; 2007.
18. Da Costa CS, Rocha NA. Sit-to-stand movement in children: a longitudinal study based on kinematics data. *Hum Mov Sci.* Aug 2013;32(4):836-846.
19. Danella A. A study of tactile preference in multiply-handicapped children. . *American Journal of Occupational Therapy.* 1973;27:457-463.

20. Davis DS, Ashby PE, McCale KL, McQuain JA, Wine JM. The effectiveness of 3 stretching techniques on hamstring flexibility using consistent stretching parameters. *J Strength Cond Res*. Feb 2005;19(1):27-32.
21. Dezateux C, Rosendahl K. Developmental dysplasia of the hip. *Lancet*. May 5 2007;369(9572):1541-1552.
22. Dietz F, Khunsree S. Idiopathic toe walking: to treat or not to treat, that is the question. *Iowa Orthop J*. 2012;32:184-188.
23. Dixon PC, Bohm H, Doderlein L. Ankle and midfoot kinetics during normal gait: a multi-segment approach. *J Biomech*. Apr 5 2012;45(6):1011-1016.
24. Eastwood DM, Menelaus MB, Dickens DR, Broughton NS, Cole WG. Idiopathic toe-walking: does treatment alter the natural history? *J Pediatr Orthop B*. Jan 2000;9(1):47-49.
25. Engelbert R, Gorter JW, Uiterwaal C, van de Putte E, Helders P. Idiopathic toe-walking in children, adolescents and young adults: a matter of local or generalised stiffness? *BMC Musculoskelet Disord*. 2011;12:61.
26. Engstrom P, Bartonek A, Tedroff K, Orefelt C, Haglund-Akerlind Y, Gutierrez-Farewik EM. Botulinum toxin A does not improve the results of cast treatment for idiopathic toe-walking: a randomized controlled trial. *J Bone Joint Surg Am*. Mar 6 2013;95(5):400-407.
27. Engstrom P, Gutierrez-Farewik E, Bartonek A, Tedroff K, Orefelt C, Haglund-Akerlind Y. Does botulinum toxin A improve the walking pattern in children with idiopathic toe-walking? *J Child Orthop* 2010;4:301-308.
28. Engstrom P, Tedroff K. The prevalence and course of idiopathic toe-walking in 5-year-old children. *Pediatrics*. Aug 2012;130(2):279-284.
29. Engstrom P, Van't Hooft I, Tedroff K. Neuropsychiatric symptoms and problems among children with idiopathic toe-walking. *J Pediatr Orthop*. Dec 2012;32(8):848-852.
30. Evans AM, Rome K, Peet L. The foot posture index, ankle lunge test, Beighton scale and the lower limb assessment score in healthy children: a reliability study. *J Foot Ankle Res*. 2012;5(1):1.
31. Evans AM, Scutter SD. Sagittal plane range of motion of the pediatric ankle joint: a reliability study. *J Am Podiatr Med Assoc*. Sep-Oct 2006;96(5):418-422.
32. Faber S. *A multisensory approach to neurorehabilitation*. In S.D. Faber (Ed). *Neurorehabilitation: A multisensory approach*. Philadelphia: Saunders; 1982.
33. Fisher A, Dunn W. Tactile defensiveness: Historial perspectives, new research- A theory grows. . *Sensory Integration Sepcial Interest Section Newsletter*. Vol 61983:1-2.
34. Flowers KR, LaStayo P. Effect of total end range time on improving passive range of motion. *J Hand Ther*. Jul-Sep 1994;7(3):150-157.
35. Fong CM, Blackburn JT, Norcross MF, McGrath M, Padua DA. Ankle-dorsiflexion range of motion and landing biomechanics. *J Athl Train*. Jan-Feb 2011;46(1):5-10.
36. Fox A, Deakin S, Pettigrew G, Paton R. Serial casting in the treatment of idiopathic toe-walkers and review of the literature. *Acta Orthop Belg*. Dec 2006;72(6):722-730.
37. Gatt A, Chockalingam N. Clinical assessment of ankle joint dorsiflexion: a review of measurement techniques. *J Am Podiatr Med Assoc*. Jan-Feb 2011;101(1):59-69.
38. Geil M, Herrin K. Is a rigid footplate as effective as an articulated AFO in controlling idiopathic toe walking? *American Academy of Orthotists & Prosthetists, 38th Academy Annual Meeting and Scientific Symposium, March 21-242012*.
39. Glennon T, Kuhaneck H, Herzberg D. The sensory processing measure-preschool (SPM-P)-part one: description of the tool and its use in the preschool environment. *Journal of Occupational Therapy, Schools, & Early Intervention*. 2011;4:42-52.
40. Goodman C, Boissonnault W, Fuller K. *Pathology: Implications for the Physical Therapist*. 2nd ed: WBSaundrrs Co; 2002.

41. Hall JE, Salter RB, Bhalla SK. Congenital short tendo calcaneus. *J Bone Joint Surg Br*. Nov 1967;49(4):695-697.
42. Harvey L, Herbert R, Crosbie J. Does stretching induce lasting increases in joint ROM? A systematic review. *Physiotherapy Research International*. 2002;7(1):1-13.
43. Hemmerich A, Brown H, Smith S, Marthandam SS, Wyss UP. Hip, knee, and ankle kinematics of high range of motion activities of daily living. *J Orthop Res*. Apr 2006;24(4):770-781.
44. Herrin K, Barner K, Geil M. Clinical outcomes after orthotic treatment of idiopathic toe walking: AFO vs. FO. *American Academy of Orthotists & Prosthetists, 38th Academy Annual Meeting and Scientific Symposium, March 21-24 2012*.
45. Hicks R, Durinick N, Gage JR. Differentiation of idiopathic toe-walking and cerebral palsy. *J Pediatr Orthop*. Mar-Apr 1988;8(2):160-163.
46. Hirsch G, Wagner B. The natural history of idiopathic toe-walking: a long-term follow-up of fourteen conservatively treated children. *Acta Paediatr*. Feb 2004;93(2):196-199.
47. Hochreiter NW, Jewell MJ, Barber L, Browne P. Effect of vibration on tactile sensitivity. *Phys Ther*. Jun 1983;63(6):934-937.
48. Iosa M, Morelli D, Nanni MV, et al. Functional taping: a promising technique for children with cerebral palsy. *Dev Med Child Neurol*. Jun 2010;52(6):587-589.
49. Irving DB, Cook JL, Young MA, Menz HB. Obesity and pronated foot type may increase the risk of chronic plantar heel pain: a matched case-control study. *BMC Musculoskelet Disord*. 2007;8:41.
50. Jacks LK, Michels DM, Smith BP, Koman LA, Shilt J. Clinical usefulness of botulinum toxin in the lower extremity. *Foot Ankle Clin*. Jun 2004;9(2):339-348.
51. Jinguji T. Toe Walking. 2012; <https://www.seattlechildrens.org/healthcare-professionals/resources/orthopedics-case-of-the-month/toe-walking-3-12/>. Accessed August, 2014.
52. Johanson M, Cuda B, Koontz J, Stell J, Abelew T. Effect of stretching on ankle and knee angles and gastrocnemius activity during the stance phase of gait. *Journal of Sports Rehabilitation*. 2009;18:521-534.
53. Kamath S, GC B. Does developmental dysplasia of the hip cause a delay in walking? *J Pediatr Orthop*. 2004;24(3):265.
54. Katz MM, Mubarak SJ. Hereditary tendo Achillis contractures. *J Pediatr Orthop*. Nov 1984;4(6):711-714.
55. Kogan M, Smith J. Simplified approach to idiopathic toe-walking. *Journal of Pediatric Orthopaedics*. 2001;21:790-791.
56. Lundequam P, Willis FB. Dynamic splinting home therapy for toe walking: a case report. *Cases J*. 2009;2:188.
57. Marcus A, Sinnott B, Bradley S, Grey I. Treatment of idiopathic toe-walking in children with autism using GaitSpot auditory speakers and simplified habit reversal. *Research in Autism Spectrum Disorders*. 2010;4:260-267.
58. Martin T, Yasukawa A. Kinesiotaping in pediatrics: Fundamentals and whole body techniques. 2003; Costa Mesa, CA.
59. Merino-Marban R, Fernandez-Rodriguez E, Lopez-Fernandez I, Mavorga-Vega D. The acute effect of kinesio taping on hamstring extensibility in university students. *Journal of Physical Education & Sport*. 2011;11(2):133-137.
60. Miller-Kuhaneck H, Henry DA, Glennon TJ, Mu K. Development of the Sensory Processing Measure-School: initial studies of reliability and validity. *Am J Occup Ther*. Mar-Apr 2007;61(2):170-175.

61. Newman CJ, Ziegler AL, Jeannet PY, Roulet-Perez E, Deonna TW. Transient dystonic toe-walking: differentiation from cerebral palsy and a rare explanation for some unexplained cases of idiopathic toe-walking. *Dev Med Child Neurol*. Feb 2006;48(2):96-102.
62. Nurse MA, Nigg BM. Quantifying a relationship between tactile and vibration sensitivity of the human foot with plantar pressure distributions during gait. *Clin Biomech (Bristol, Avon)*. Nov 1999;14(9):667-672.
63. Oetgen ME, Peden S. Idiopathic toe walking. *J Am Acad Orthop Surg*. May 2012;20(5):292-300.
64. O'Sullivan K, Murray E, Sainsbury D. The effect of warm-up, static stretching and dynamic stretching on hamstring flexibility in previously injured subjects. *BMC Musculoskelet Disord*. 2009;10:37.
65. Parham L, Ecker C, Miller-Kuhaneck H, Henry DA, Glennon TJ. *Sensory Processing Measure (SPM) Manual*. Los Angeles: Western Psychological Services; 2007.
66. Pernet J, Billiaux A, Auvin S, et al. Early onset toe-walking in toddlers: a cause for concern? *J Pediatr*. Sep 2010;157(3):496-498.
67. Pfeiffer B, Kinnealey M. Treatment of sensory defensiveness in adults. *Occup Ther Int*. 2003;10(3):175-184.
68. Protopapadaki A, Drechsler WI, Cramp MC, Coutts FJ, Scott OM. Hip, knee, ankle kinematics and kinetics during stair ascent and descent in healthy young individuals. *Clin Biomech (Bristol, Avon)*. Feb 2007;22(2):203-210.
69. Rabin A, Kozol Z. Weightbearing and nonweightbearing ankle dorsiflexion range of motion: are we measuring the same thing? *J Am Podiatr Med Assoc*. Sep-Oct 2012;102(5):406-411.
70. Radford J, Burns J, Buchbinder R, Landorf K, Cook C. Does stretching increase ankle dorsiflexion range of motion? A systematic review. *Br J Sports Med*. 2006;40:870-875.
71. Reid DA, McNair PJ. Passive force, angle, and stiffness changes after stretching of hamstring muscles. *Med Sci Sports Exerc*. Nov 2004;36(11):1944-1948.
72. Ribot-Ciscar E, Vedel JP, Roll JP. Vibration sensitivity of slowly and rapidly adapting cutaneous mechanoreceptors in the human foot and leg. *Neurosci Lett*. Sep 25 1989;104(1-2):130-135.
73. Sala DA, Shulman LH, Kennedy RF, Grant AD, Chu ML. Idiopathic toe-walking: a review. *Dev Med Child Neurol*. Dec 1999;41(12):846-848.
74. Santonja Medina FM, Sainz De Baranda Andujar P, Rodriguez Gracia PL, Lopez Minarro PA, Canteras Jordana M. Effects of frequency of static stretching on straight leg raise in elementary school children. *Journal of Sports Medicine and Physical Fitness*. 2007;47(3):304-308.
75. Schoenfeld BJ. Squatting kinematics and kinetics and their application to exercise performance. *J Strength Cond Res*. Dec 2010;24(12):3497-3506.
76. Segal R, Beyer C. Integration and application of a home treatment program: a study of parents and occupational therapists. *Am J Occup Ther*. Sep-Oct 2006;60(5):500-510.
77. Sobel E, Caselli MA, Velez Z. Effect of persistent toe walking on ankle equinus. Analysis of 60 idiopathic toe walkers. *J Am Podiatr Med Assoc*. Jan 1997;87(1):17-22.
78. Solan MC, Kohls-Gatzoulis J, Stephens MM. Idiopathic toe walking and contractures of the triceps surae. *Foot Ankle Clin*. Jun 2010;15(2):297-307.
79. Soucie JM, Wang C, Forsyth A, et al. Range of motion measurements: reference values and a database for comparison studies. *Haemophilia*. May 2011;17(3):500-507.
80. Stott NS, Walt SE, Lobb GA, Reynolds N, Nicol RO. Treatment for idiopathic toe-walking: results at skeletal maturity. *J Pediatr Orthop*. Jan-Feb 2004;24(1):63-69.
81. Stricker SJ, Angulo JC. Idiopathic toe walking: a comparison of treatment methods. *J Pediatr Orthop*. May-Jun 1998;18(3):289-293.
82. Tabrizi P, McIntyre WM, Quesnel MB, Howard AW. Limited dorsiflexion predisposes to injuries of the ankle in children. *J Bone Joint Surg Br*. Nov 2000;82(8):1103-1106.

83. van Bemmel AF, van de Graaf VA, van den Bekerom MP, Vergroesen DA. Outcome after conservative and operative treatment of children with idiopathic toe walking: a systematic review of literature. *Musculoskelet Surg.* Aug 2014;98(2):87-93.
84. Vander Linden DW, Wilhelm IJ. Electromyographic and cinematographic analysis of movement from a kneeling to a standing position in healthy 5- to 7-year-old children. *Phys Ther.* Jan 1991;71(1):3-15.
85. Weeks S, Boshoff K, Stewart H. Systematic review of the effectiveness of the Wilbarger protocol with children. *Pediatric Health, Medicine, and Therapeutics.* 2012;3:79-89.
86. Wegener C, Hunt AE, Vanwanseele B, Burns J, Smith RM. Effect of children's shoes on gait: a systematic review and meta-analysis. *J Foot Ankle Res.* 2011;4:3.
87. Williams CM, Tinley P, Curtin M. Idiopathic toe walking and sensory processing dysfunction. *J Foot Ankle Res.* 2010;3:16.
88. Williams CM, Tinley P, Curtin M. The Toe Walking Tool: a novel method for assessing idiopathic toe walking children. *Gait Posture.* Oct 2010;32(4):508-511.
89. Williams CM, Tinley P, Curtin M, Nielsen S. Vibration perception thresholds in children with idiopathic toe walking gait. *J Child Neurol.* Aug 2012;27(8):1017-1021.
90. Williams CM, Tinley P, Curtin M, Wakefield S, Nielsen S. Is idiopathic toe walking really idiopathic? The motor skills and sensory processing abilities associated with idiopathic toe walking gait. *J Child Neurol.* 2013.
91. Williams CM, Tinley P, Rawicki B. Idiopathic toe-walking: have we progressed in our knowledge of the causality and treatment of this gait type? *J Am Podiatr Med Assoc.* May 2014;104(3):253-262.
92. Young R, Nix S, Wholohan A, Bradhurst R, Reed L. Interventions for increasing ankle joint dorsiflexion: a systematic review and meta-analysis. *J Foot Ankle Res.* 2013;6(1):46.